

CLAIM

1. A monoclonal antibody which binds to a glycoprotein antigen that has a molecular weight of 200 kD or more (SDS-PAGE), exists in human lung adenocarcinoma cells and is secreted by human lung adenocarcinoma.

2. The monoclonal antibody according to claim 1 wherein it reacts with MAA lectin and PNA lectin but does not react with GNA lectin, SNA lectin and DSA lectin.

3. The monoclonal antibody according to claim 1 or 2 wherein isotype of the aforementioned antibody is IgM.

4. The monoclonal antibody according to any one of claims 1 to 3 wherein it is produced by a hybridoma cell strain TRD-L1, TRD-L2 or TRD-L3 obtained by the fusion between mouse myeloma cells and spleen cells of a mouse immunized with a human lung adenocarcinoma cell secretion component.

5. The monoclonal antibody according to any one of claims 1 to 4 wherein the aforementioned antibody is an Fab, F(ab)₂ or Fv fragment.

6. A glycoprotein antigen which is a glycoprotein having a molecular weight of 200 kD or more (SDS-PAGE), exists in cells of human lung adenocarcinoma and is secreted by lung adenocarcinoma.

7. The glycoprotein antigen according to claim 6 wherein it reacts with MAA lectin and PNA lectin but does not react with GNA lectin, SNA lectin and DSA lectin.

09350899.01299
662720 6680560

8. An immunoassay method for cancer diagnosis use which comprises using the monoclonal antibody of any one of claims 1 to 5.

9. The immunoassay method for cancer diagnosis use according to claim 8 wherein the glycoprotein antigen of claim 6 or 7 is measured using the monoclonal antibody of any one of claims 1 to 5.

09350899 071299
662720 66805660

ADD
A1